The Hidden Impact of Molluscum Contagiosum: Survey of Caregivers' Experiences with Diagnosis and Management

Pearl Kwong¹, Adelaide Hebert², Collette Utley³, Melissa Olivadoti⁴ ¹Solutions Through Advanced Research, Jacksonville, FL; ² UTHealth McGovern Medical School, Houston, TX; ³Gold Skin Care Center, Nashville, TN; ⁴Verrica Pharmaceuticals Inc., West Chester, PA

INTRODUCTION

- Molluscum contagiosum ("molluscum") is a common pediatric viral skin infection that is considered benign and self-limiting. However, it may affect the quality of life in patients and their caregivers dealing with the disease.
- The experience of caregiving for a child with molluscum is largely unknown, including the emotional burden, effort for diagnosis, and treatment modalities used for molluscum.

METHODS

- Parents, caregivers, and legal guardians (ages 18+, 80% female, 20% male) of children ages 3–16 years diagnosed with molluscum in the past 4 years were recruited from a panel of previously recruited respondents to answer a 15-minute paid online survey.
- Question topics included the type of health care professionals (HCPs) consulted about diagnosis, treatment (both in the clinic and at home), and the impact of the virus on the caregiver and the child.
- The margin of error at the 95% confidence interval for the total sample was ± 7.7%.

RESULTS

	Respondents (N=154)
Household Information	
Median number of people in the home	4
Median age of children in the home in years	8
Household size – No. (%)	
2-3 People	44 (28)
4-5 People	94 (61)
6+ People	16 (10)
Age of Children in the Home – No. (%)	
Under 3 years	48 (31)
4-8 years	80 (52)
9-12 years	77 (50)
13-16 years	67 (43)
Caregiver Gender – No. (%)	
Male	31 (20)
Female	123 (80)
Molluscum Status – No. (%)	
Child currently has molluscum	116 (75)
Child had molluscum in the last 4 years	39 (25)
Spread of molluscum from one child to another in the home*	65 (60)
Ethnicity – No. (%)	
White/Caucasian	131 (85)
Black/African American	16 (10)
Asian	6 (4)
Native Hawaiian/Pacific Islander	1 (1)
American Indian/Alaskan Native	2 (1)
Other	3 (2)
Homestead information – No. (%)	
City	79 (51)
Suburb	65 (42)
Small town/rural	10 (7)

* In homes of >1 child, n=108.




Figure 2. Types of HCPs Consulted About Molluscum Contagiosum




Figure 3. HCP Treatment Recommendations





Figure 4. "Which of the following do you remember the doctor telling you?"*



The average number of HCPs consulted per caregiver was 1.95.




Figure 5. "How much of an impact has molluscum had on your child's life (right figure) or on your life (left figure)?





Figure 6. "Which of the following ways of dealing with molluscum did you decide on for your child?"





- These survey results indicate that patients with molluscum receive a diagnosis from numerous sources and that many molluscum patients do not receive treatment.
- Caregivers may use more than one treatment modality to rid their children of the infection and are likely to use at-home remedies, including physical disruption of lesions.
- Spread to another child in the household was common despite despite being told about the infectious nature of the virus.
- Caregivers reported a substantial impact on their quality of life and on the activities of their children with molluscum.
- There is an unmet need for education of HCPs regarding the diagnosis and impact of molluscum, in addition to safe and effective treatment options for the infection.

Disclosures

The studies were sponsored by Verrica Pharmaceuticals Inc. Editorial support was provided by Versant Learning Solutions and funded by Verrica Pharmaceuticals Inc. The authors have received the following from Verrica Pharmaceuticals: P. Kwong: H, C; A. Herbert: H. C. Utley: H; M. Olivadoti: E H=honoraria; C=clinical funds; S=stocks; E=employee.

